



Manuscript version: Author's Accepted Manuscript

The version presented in WRAP is the author's accepted manuscript and may differ from the published version or Version of Record.

Persistent WRAP URL:

<http://wrap.warwick.ac.uk/112565>

How to cite:

Please refer to published version for the most recent bibliographic citation information. If a published version is known of, the repository item page linked to above, will contain details on accessing it.

Copyright and reuse:

The Warwick Research Archive Portal (WRAP) makes this work by researchers of the University of Warwick available open access under the following conditions.

Copyright © and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable the material made available in WRAP has been checked for eligibility before being made available.

Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

Publisher's statement:

Please refer to the repository item page, publisher's statement section, for further information.

For more information, please contact the WRAP Team at: wrap@warwick.ac.uk.

Management of first-trimester miscarriage: a systematic review and network meta-analysis

Bassel H.Al Wattar^{1,2}, Nilaani Murugesu¹, Aurelio Tobias³, Javier Zamora^{1,4}, Khalid S. Khan^{1, 5}

¹Women's Health Research Unit, Barts and the London School of Medicine and Dentistry, Queen Mary University London, London, UK

²Warwick Medical School, University of Warwick, Coventry, UK

³Institute of Environmental Assessment and Water Research, Spanish Council for Scientific Research (CSIC), Barcelona, Spain

⁴Clinical Biostatistics Unit, Ramon y Cajal Hospital (IRYCIS) and CIBER Epidemiology and Public Health, Madrid, Spain

⁵Multidisciplinary Evidence Synthesis Hub (mEsh), Barts and the London School of Medicine and Dentistry, Queen Mary University of London, London, UK

Corresponding Author:

Dr. Bassel H.Al Wattar

Barts and the London School of Medicine and Dentistry, Queen Mary University London, London E1 2DD

E-Mail: b.wattar@qmul.ac.uk

Running title: Management of first-trimester miscarriage

25	Table of content
26	Introduction
27	Methods
28	Search strategy
29	Selection criteria and data extraction
30	Primary and secondary outcomes
31	Types of treatment for first trimester miscarriage
32	Quality assessment of risk of bias
33	Statistical analysis
34	Patient involvement
35	Results
36	Characteristics of included studies
37	Risk of bias
38	Primary outcome
39	Secondary outcomes
40	Discussion
41	Main findings
42	Strength and limitations
43	Interpretation of findings
44	Conclusions
45	

Abstract

Background: First-trimester miscarriage affects up to a quarter of women worldwide. With many competing treatment options available, there is a need for a comprehensive evidence synthesis.

Objectives and rationale: We conducted a systematic review and network meta-analysis to assess the effectiveness and safety of treatment options for first-trimester miscarriage: expectant management (EXP), sharp dilation and curettage (D+C), electric vacuum aspiration (EVAC), manual vacuum aspiration (MVA), misoprostol alone (MISO), mifepristone+misoprostol (MIFE+MISO) and misoprostol plus electric vacuum aspiration (MISO+EVAC).

Search methods: We searched MEDLINE, Embase, CINAHL, AMED and Cochrane Library from inception till June 2018. We included randomised trials of women with first-trimester miscarriage (<14 weeks gestation) and conducted a network meta-analysis generating both direct and mixed evidence on the effectiveness and side effects of available treatment options. The primary outcome was complete evacuation of products of conception. We assessed the risk of bias and the global network inconsistency. We compared the surface under the cumulative ranking curve (SUCRA) for each treatment.

Outcomes: A total of 46 trials (9250 women) were included. The quality of included studies was overall moderate with some studies demonstrating a high risk of bias. We detected unexplained inconsistency in evidence loops involving MIFE+MISO and adjusted for it. EXP had lower effectiveness compared to other treatment options. The effectiveness of medical treatments was similar compared to surgery. Mixed evidence of low confidence suggests increased effectiveness for MIFE+MISO compared to MISO alone (RR 1.49, 95% CI 1.09-2.03). Side effects were similar among all options. Fewer women needed analgesia following EVAC compared to MISO (RR for MISO 0.43, 95% CI 0.27-0.68) and in the EXP group

71 compared to EVAC (RR 2.07, 95% CI 1.25-3.41). MVA had higher ranking (low likelihood)
72 for post-treatment infection and serious complications (SUCRA 87.6%, 79.2% respectively)
73 with the highest likelihood for post-treatment satisfaction (SUCRA 98%).

74 **Wider implications:** Medical treatments for first-trimester miscarriage have similar
75 effectiveness and side effects compared to surgery. The addition of MIFE could increase the
76 effectiveness of MISO and reduce side effects, although evidence is limited due to
77 inconsistency. EXP has lower effectiveness compared to other treatment options.

78
79 **Systematic review registration:** Prospero CRD42016048920

80
81 **Keywords:** miscarriage, pregnancy loss, first trimester, effectiveness, woman, systematic
82 review, network meta-analysis.

Introduction

First trimester miscarriage, the most common time of pregnancy loss, is estimated to affect up to a quarter of pregnant women in their lifetime (Wang *et al.*, 2003). Miscarriage can lead to significant clinical and emotional morbidity, affecting the couples' quality of life (Jurkovic *et al.*, 2013). Providing patient-centred care can help to reduce the psychological sequelae associated with miscarriage (van den Berg *et al.*, 2017) such as increased anxiety, depression, grief and low self-esteem (Frost and Condon, 1996; Swanson *et al.*, 2009). The burden of miscarriage on healthcare resources is significant, leading to over 50,000 hospital admissions annually in the UK (The National Institute for Health and Care Excellence, 2012), with a similar impact in other developed countries (Queensland Clinical Guidelines, 2015; The American College of Obstetricians and Gynecologists, 2015).

Various treatment options exist for couples experiencing first-trimester miscarriage; these are broadly categorised into expectant, medical and surgical groups (Trinder *et al.*, 2006). The wide use of less invasive treatments such as prostaglandins and manual vacuum evacuation could reduce the need for surgical interventions under general anaesthesia and the number of hospital admissions (Jurkovic *et al.*, 2013; Sotiriadis *et al.*, 2005). Misoprostol is currently the most used drug for treating miscarriage, however, there is no consensus on the best dose and route of its administration (Neilson *et al.*, 2013). Combining medical and surgical treatments is common, though evidence to support this practice is imprecise (Fang *et al.*, 2009). Evidence concerning the effectiveness and safety of available treatment options is limited to pairwise comparisons in randomised trials and their meta-analyses (Nanda *et al.*, 2006; Neilson *et al.*, 2013; Sotiriadis *et al.*, 2005; Tunçalp *et al.*, 2010).

There is a need for a comprehensive evidence synthesis to compare the effectiveness and safety of the available treatment options. We conducted a systematic review and a network meta-analysis of randomised trials (comparing different treatments for a particular condition using the estimated effect size from direct and indirect comparisons) (Al Wattar *et al.*, 2017) to assess the effectiveness and side effects of available treatment options for complete evacuation of products of conception in women experiencing first-trimester miscarriage.

Methods

We conducted our systematic review according to a prospectively registered protocol (Prospero CRD42016048920) and reported the findings to comply with the extended PRISMA guidelines (Hutton *et al.*, 2015). The final author affirms that the manuscript is an honest, accurate, and transparent account of the study being reported; no important aspects of the study have been omitted; and there are no discrepancies from the planned study protocol.

Search strategy

We searched the following electronic databases for randomised trials comparing any treatment option for first-trimester miscarriage from inception until June 2018 (MEDLINE, Embase, CINAHL, AMED and Cochrane Library). We developed a multi-step search strategy and adjusting it appropriately for each database (not shown). No search filters were applied. We conducted supplementary searches in Google Scholar and Scopus. We manually screened bibliographies of reviewed articles to identify any additional relevant trials. Articles in non-English language were obtained and translated if deemed relevant. We contacted authors for further information when needed, but no unpublished data were included. We reviewed all available systematic reviews on the management of first-trimester miscarriage to identify any additional studies.

Selection criteria and data extraction

We included all randomised trials that evaluated any treatment option in women with first-trimester miscarriage (defined as a spontaneous loss of a non-viable intrauterine pregnancy between 0 and 14 weeks' gestation) (The National Institute for Health and Care Excellence, 2012). Studies that included a combination of two treatment options (e.g. medical plus surgical) were included. Studies with multiple comparison arms were also included. We excluded quasi-randomised studies and those reporting on elective termination of pregnancy. Studies that compared variations of the same treatment in both arms (e.g. misoprostol 400 µg vs misoprostol 600 µg) were reported narratively and excluded from the meta-analysis. Studies that reported on secondary outcomes only were also excluded. We manually extracted data, using a bespoke electronic tool, on the place of the study, the publication journal, treatment settings, population characteristics, the treatment options evaluated, including its dose and route where applicable, and primary and secondary outcomes. The selection and data extraction processes were conducted in duplicate by two independent reviewers (BHA and NM). Any disagreement was resolved by discussion with a third reviewer (KSK).

Primary and secondary outcomes

Our primary outcome was complete evacuation of products of conception, defined clinically or on ultrasound as an empty uterine cavity without the need for further treatment. Secondary outcomes were: serious complications (defined as a composite of any of the following: uterine perforation, cervical tear, hysterectomy, laparotomy, Asherman's syndrome, and death), need for blood transfusion, post-treatment infection/pelvic inflammatory disease, nausea, vomiting, diarrhoea, fever (>38 °C with no evidence of infection), patient

satisfaction, mean hospital stay (days), visual analogue pain scores, anxiety, depression and need for analgesia.

Types of treatment for first trimester miscarriage

Treatment options were grouped into five categories: expectant (defined as conservative management with no active intervention including placebo), medical (defined as any medical drug of any dose, route and format to achieve uterine evacuation), placebo (defined as a planned placebo intervention within a trial settings), surgical (defined as any surgical instruments used under general or local anaesthesia to achieve uterine evacuation) and a combination of any medical plus surgical treatment used consecutively. To reduce inconsistency in the network, we combined conservative and placebo treatments under the same label (expectant management). We also excluded uncommon medical drugs that were reported in single studies (e.g. Methotrexate) and reported on them narratively.

Quality assessment of risk of bias

We assessed the risk of bias in all included studies in duplicate by two independent reviewers (BHA and NM) using the Cochrane risk of bias assessment tool (Higgins *et al.*, 2011). This included assessment of the following items: randomisation and sequence generation, allocation concealment, blinding and performance, outcome assessment, completeness of outcome data and selective outcome reporting. Unblinded studies were not penalised in the risk of bias assessment due to the nature of the treatments that makes blinding non-feasible. Quality assessment was performed in duplicate by two independent reviewers.

Statistical analysis

182 We performed standard pairwise meta-analyses using a random-effects model (Sutton *et al.*,
 183 2000) and network meta-analysis within a frequentist framework with multivariate meta-
 184 analysis models (White *et al.*, 2012), exploiting the direct and indirect randomised evidence
 185 to determine the relative effects and ranking. We reported on direct evidence (from head to
 186 head comparison of treatments) and mixed evidence (combining both direct and indirect
 187 evidence from comparison of treatments) using weighted mean difference (WMD) for
 188 continuous outcomes and risk ratios (RR) for dichotomous outcomes with 95% confidence
 189 intervals (CI). We also computed the probability that each treatment is the most effective, as
 190 well as the surface under the cumulative ranking curve (SUCRA) to compare the relative
 191 ranking probability of each treatment (Chaimani *et al.*, 2013; Salanti *et al.*, 2011). Providing
 192 a cumulative rankogram adjusts for any uncertainty in the relative treatment effect where
 193 limited evidence exists (Chaimani *et al.*, 2013). A cumulative rank provides the probability
 194 for each treatment to be the best among the rang of available treatment options; the SUCRA
 195 is a transformation of the mean rank accounting for the location and variance of available
 196 treatment effects to generate a treatment hierarchy (Salanti *et al.*, 2011).
 197 In pairwise meta-analyses, we estimated different heterogeneity variances for each pairwise
 198 comparison, using the I^2 index, to capture the percentage of variation that is not due to
 199 chance. In the network meta-analysis, we assumed a common estimate for the heterogeneity
 200 variance across the different comparisons. To check the assumption of consistency in the
 201 entire network, we used the design-by-treatment model (Higgins *et al.*, 2012). In case of
 202 whole network inconsistency, we investigated differences between direct and indirect
 203 evidence using the loop-specific approach (Bucher *et al.*, 1997), assuming a common
 204 heterogeneity estimate within each loop (a loop of evidence exist when numerous trials
 205 compare a minimum of three treatments e.g A vs B vs C) (Veroniki *et al.*, 2013). We
 206 investigated any detected inconsistency and adjusted for unexplained inconsistency within the

network using established models in STATA (Riley *et al.*, 2017). All analyses were done using STATA statistical software, release 14 (StataCorp, College Station, TX, 2015).(Chaimani *et al.*, 2013; White, 2011; White *et al.*, 2012).

Patient involvement

We did not involve a patient representative in the design of our study. We consulted the James Lind Library and previous Cochrane reviews to identify the primary outcome and other outcomes of interest to stakeholders.

Results

Characteristics of included studies

Our electronic search identified 3648 potentially relevant studies. Of these, we excluded 3523 after reviewing titles and abstracts. The remaining 125 studies were assessed in full. Eleven studies were identified from screening bibliographies and were assessed in full. We excluded 90 studies: five reporting on the use of methotrexate, dinoprestone, mifepristone alone, laminaria, gemeprost (Autry *et al.*, 1999; Al Inizi and Ezimokhai, 2003; Johnson *et al.*, 1997; Lelaidier *et al.*, 1993), 20 comparing different dosages, routes or formats of misoprostol against each other, 15 non- or quasi-randomised, 16 not meeting the inclusion criteria and 34 not reporting the primary outcome. In total 46 randomised trials reporting on 9250 women were included, of these two were in Portuguese (Holanda *et al.*, 2003; Pereira *et al.*, 2006) and one in Norwegian (Karlsen, Jørn-Hugo; Hjalmar, 2001) (Figure 1).

A third of included trials were conducted in European countries (14/46, 30.4%) and fourteen in Asian countries (14/46, 30.4%). Most studies included a two arms comparison and four included three arms. The median study sample size was 60 (range 12-402). The majority of trials were conducted in tertiary healthcare settings (35/46, 76.1%). One study was conducted

in outpatient settings. Eight were multicentre randomised trials (8/46, 17.3%). Table I provides a summary of the characteristics of included trials.

Risk of bias

The quality of included studies was overall moderate with some studies demonstrating a high risk of bias (Supplementary Figure S1). Nine studies had a high risk of bias for randomisation (9/46, 19.5%) and ten (10/46, 21.7%) had a high risk of bias for allocation concealment. Outcomes assessment (i.e. attrition) was judged to have a high risk of bias in six studies, and was inadequate in 15 studies (15/46, 32.6%) but good in 25 studies (25/46, 60.9%). Six studies had a high risk of bias for detection (i.e. selective reporting) (6/46, 13%) and 14 had a high risk of bias in outcomes reporting (i.e. incomplete data) (14/46, 30.4%). Conflict of interest was declared as not present in only seven studies (7/46, 15.2%) and was not reported on in the remaining studies. Only four studies were double blinded and these were studies comparing medical treatments to placebo (3/46, 6%) (Bagratee *et al.*, 2004; Blohm *et al.*, 2005; Lister *et al.*, 2005; Sinha *et al.*, 2018). A summary of risk of bias assessment on included trials is provided in Supplementary Table SII.

Primary outcome

Our network for the primary outcome included 46 randomised trials (9250 women) comparing seven treatment options: expectant management (EXP)(19 trials, 1587 women), sharp dilation and curettage (D+C)(5 trials, 247 women) , electric vacuum aspiration under general anaesthesia (EVAC)(19 trials, 1766 women), manual vacuum aspiration under local anaesthesia (MVA)(12 trials, 1671 women), misoprostol alone (MISO)(32 trials, 3017 women), mifepristone + misoprostol (MIFE+MISO)(9 trials, 932 women), sequential

256 misoprostol + electric vacuum aspiration under general anaesthesia (MISO+EVAC)(1 trial,
 257 30 women) (Figure 2).

258 Both direct and mixed evidence supported the overall inferiority of EXP compared to most
 259 treatment options for achieving complete evacuation of products of conception (EXP vs
 260 MISO RR 0.76, 95% CI 0.65-0.89; EXP vs EVAC RR 0.68, 95% CI 0.59-0.79; EXP vs D+C
 261 RR 0.73, 95% CI 0.57-0.94; MISO+EVAC vs EXP RR 1.35, 95% CI 1.10-1.66; MVA vs
 262 EXP RR 1.46, 95% CI 1.19-1.79) (Figure 3). All surgical treatments (MVA, EVAC and
 263 D+C) demonstrated similar effectiveness for achieving the primary outcome. This was also
 264 the case when comparing MISO against each of the surgical treatment options (MVA vs
 265 MISO RR 1.10, 95% CI 0.92-1.33; EVAC vs MISO RR 1.11, 95% CI 0.97-1.27; D+C vs
 266 MISO RR 1.03, 95% CI 0.82-1.30). Direct evidence on the use of MISO+EVAC was drawn
 267 from one trial only (MISO+EVAC vs MISO, RR 2.86, 95% CI 1.45-5.64; data not shown)
 268 and mixed evidence supports its superiority only over EXP (RR 1.35, 95% CI 1.10-1.66).
 269 Mixed evidence did not support the use of MIFE+MISO compared to using MISO alone to
 270 increase effectiveness (RR 1.43, 95% CI 0.87-2.36). However, we detected significant
 271 inconsistency between direct and mixed evidence for MISO vs MISO+EVAC; EVAC vs
 272 MISO; EVAC vs EXP; EVAC vs MIFE+MISO and EXP vs MISO (Supplementary Table
 273 SI). The overall by network inconsistency analysis was significant at $p=0.003$. Adjusting for
 274 inconsistency, mixed evidence favoured the addition of MIFE to MISO to improve
 275 effectiveness (MIFE+MISO vs MISO RR 1.49, 95% CI 1.09-2.03) in contrast to
 276 MISO+EVAC vs MISO (RR 0.63, 95% CI 0.51-0.79) (Supplementary Figure S3).
 277

278 The surface under the cumulative ranking curve for treatment effectiveness was highest for
 279 MIFE+MISO (SUCRA 89.3%) followed by EVAC (SUCRA 76.2%). EXP was ranked as the
 280 least effective treatment (SUCRA 24%) (Figure 4). Visual analysis of our funnel plot

demonstrates a reasonable distribution of effect size with limited evidence of small study effect (Supplementary Figure S4).

Secondary outcomes

Meta-analysis of mixed evidence demonstrated no difference for any of the following outcomes between medical and surgical treatment options: need for blood transfusion, post-treatment infection, serious complications, diarrhoea, vomiting, nausea and fever (Supplementary Figures S5-11). Compared to MISO, MIFE+MISO was associated with a lower risk ratio for developing fever (RR 0.33, 95% CI 0.19-0.57), nausea (RR 0.42, 95% CI 0.24-0.72) and vomiting (RR 0.55, 95% CI 0.32-0.94). Fewer women needed analgesia post treatment in the EVAC group compared to MISO (RR 0.43, 95% CI 0.27-0.68). Those who opted for EXP also used more analgesia compared to EVAC (RR 2.07, 95% CI 1.25-3.41) (Supplementary Figure S12). Women's satisfaction was similar for all the treatment options (Supplementary Figure S13). Supplementary Table SIII provides a summary of effect estimates for all secondary outcomes across treatment options.

Table II summaries the calculated SUCRA and mean rank for the secondary outcomes by the treatment options. Generally, MIFE+MISO had high ranking (low likelihood) for causing common gastrointestinal (GI) side effects (nausea (SUCRA 93.1%), vomiting (SUCRA 84%) and diarrhoea (SUCRA 63.2%)) and fever (SUCRA 86.8%). MVA had higher ranking (low likelihood) for post-treatment infection (SUCRA 87.6%) and serious complications (SUCRA 79.2%) with the highest likelihood for post-treatment satisfaction (SUCRA 98%). Women opting for EVAC had higher likelihood of requiring post-treatment blood transfusion (SUCRA 14.7%).

Discussion

Main findings

Our comprehensive meta-analysis showed that for managing first-trimester miscarriage, EXP had lower effectiveness to achieve complete evacuation of products of conception compared to other treatment options. Overall, there was similar effectiveness for the medical (MIFE+MISO and MISO) and the surgical options (MVA, D+C, and EVAC), with similar safety profiles reported. There was limited evidence to support the use of MISO+EVAC with no information on its safety profile. Evidence on the use of MIFE+MISO suffered from significant inconsistency. Overall, the addition of MIFE to MISO seems to improve its effectiveness with reduced likelihood of side effects but more research is needed to address the perceived inconsistency between direct and indirect evidence. Women's satisfaction was similar for all the options compared.

Currently, EXP is recommended as the first-line treatment option for first-trimester miscarriage (The National Institute for Health and Care Excellence, 2012). Women opting for this approach should be counselled objectively about the chances of needing further treatment, potential complications such as requiring blood transfusion (SUCRA 36.3%) or more analgesia (SUCRA 37.4%), and the availability of other effective treatment options. Excessive bleeding and repeated blood transfusion contribute to prolonged hospital stays and long-term adverse outcomes such as alloimmunisation (Royal College of Obstetricians and Gynaecologists, 2015) which are infrequently assessed in randomised trials.

Strength and limitations

This review, to our knowledge, is the first to provide a comprehensive evidence synthesis with network meta-analysis on all current treatment options for first-trimester miscarriage.

We conducted a systematic review of the literature with no search limitations. We assessed and found little evidence of small study effect with the funnel plot analysis raising confidence in our findings. We assessed the risk of bias using the Cochrane risk of bias assessment tool (Higgins *et al.*, 2011) which demonstrated low to moderate risk of bias in the majority of included studies. Compared to previously conducted meta-analysis (Nanda *et al.*, 2006; Neilson *et al.*, 2006, 2013; Sotiriadis *et al.*, 2005; Wen *et al.*, 2008), our study provides higher confidence supporting the role of medical treatment options for first trimester miscarriage, incorporating indirect evidence and ranking treatments likelihood for effectiveness and side effects.

Our findings are not without limitations. We were unable to accommodate for potential effect modifiers such as variation in population characteristics relevant to age, parity, size of products of conception, presence of side effects before randomisation and treatment settings. A large gestation sac might require a higher doses of MISO to achieve complete evacuation (Neilson *et al.*, 2006). Evidence on some treatment options, such as MVA, was sought primarily from low/middle income countries, which could suggest variations in local practice and geographical bias to one treatment option over the others.

There were variations in the ultrasound criteria used to diagnose the type of miscarriage (missed vs incomplete) and the primary outcome of complete evacuation of products of conception. The use of a standardised ultrasound criteria for the diagnosis of miscarriage is only recent and some of the included trials pre-date the currently established guidelines (The National Institute for Health and Care Excellence, 2012). To be pragmatic, we opted to keep those trials and offer a comprehensive and accurate review of the available literature. Similarly, there was variation in the type of included miscarriages (missed vs incomplete) in

each trial with some trials randomising either or both or simply not reporting on it (Table 1). Due to the risk of inconsistency, we were unable to generate evidence on the management of each type of miscarriage and our findings remain pragmatic. Such variation could be best addressed using an individual participant data meta-analysis.

There was inconsistency within the network (Supplementary Table SI) specifically within evidence loops comparing MIFE+MISO to other treatment options. We were unable to attribute this inconsistency to a particular effect modifier and adjusted for it using established models. Inconsistency could be attributed to the variations in the dosages and the routes of administration of MISO among included trials. Typically, MISO is used in sequential doses of 200 mcg and stopped once complete evacuation of products is achieved; this could present inherent inconsistency among trials. Quality evidence on the most effective dose with the least side effect is yet to emerge (Neilson *et al.*, 2006).

Variations in defining endpoints and the follow-up period limited the information on important long term outcomes such as uterine adhesions, pre-term birth and future fertility. Recent evidence suggest an increased risk for pre-term birth with multiple dilation and curettage (Lemmers *et al.*, 2015). Future work should focus on following up randomised cohorts to capture such outcomes.

To be pragmatic, evidence on MISO+EVAC, sought from one trial (Fang *et al.*, 2009), was kept within our network in view of its wide use in current practice. The findings of this trial should be interpreted with caution due to its small sample size, moderate risk of bias and limited reporting on secondary outcomes. We planned to report on four additional outcomes

in our protocol (hospital stay, changes in haemoglobin, anxiety, and depression). This, however, was not possible due to the large variability in reported end points. We judged blinding to be possible in seven studies (Bagratee *et al.*, 2004; Blohm *et al.*, 2005; Herabutya and O-Prasertsawat, 1997; Lister *et al.*, 2005; Ngai *et al.*, 2001; Nielsen *et al.*, 1999; Wood and Brain, 2002). Of these, only four (Bagratee *et al.*, 2004; Blohm *et al.*, 2005; Lister *et al.*, 2005; Sinha *et al.*, 2018) were blinded, introducing a potential risk of bias. Lack on information on blinding for outcomes assessment is another limitation in the included studies.

Interpretation of findings

Our study supports the use of medical treatments as a potential substitute for surgery, however, studies to establish the lowest effective dose of MISO are needed (Neilson *et al.*, 2006). A higher dose of MISO is likely to cause more side effects such as nausea and vomiting (Tang *et al.*, 2007). Medical management could be considered as a cost-effective first-line treatment option. The woman's preference is an important factor to consider when offering the various treatment options, often influenced by their carer's advice. There was seldom consideration in the included studies for reporting outcomes important to the women undergoing miscarriage, such as post-treatment anxiety and depression. None of the included studies reported on the tolerability of each treatment option which can aid women to identify their preferred choice. Developing a core outcome set with input from all stakeholders should be considered in future research (Khan, 2016).

Recently, MIFE has been more commonly combined with MISO to improve the effectiveness of medical treatment for uterine evacuation (Spitz *et al.*, 1998). Our analysis, seeking direct and mixed evidence, suggests some added value compared to using MISO alone for first-

trimester miscarriage but with limited confidence due to the perceived inconsistency among included trials. Considering its high cost, a cost-effectiveness evaluation is needed to establish the value of using MIFE+MISO routinely. Using MISO for priming the cervix before EVAC has been suggested to reduce the need for dilation and trauma to the endometrium (Lawrie *et al.*, 1996). Evidence to support the effectiveness and safety of this practice for managing first trimester miscarriage is scarce (only one randomised trial of 75 women) (Fang *et al.*, 2009) and more trials are needed to justify the potentially increased cost and side effects.

Outpatient use of MVA with direct access to operating theatres could offer cost reduction (Magotti *et al.*, 1995). While EXP is arguably cheaper than other treatment options, the higher probability of complications might increase its associated cost. There is a need for a comprehensive economic evaluation with extended decision models to accommodate for the effectiveness of all available treatment options and potential adverse outcomes (Strand, 2015). Comprehensive policymaking including all available treatment options could offer better value for money and facilitate higher patient satisfaction (Dalton *et al.*, 2015; Molnar *et al.*, 2000; Wallace *et al.*, 2010) (Supplementary Figure S2).

Our study provides important insight for various stakeholders involved in caring for women with first trimester miscarriage. Future work should aim to involve stakeholders' views prospectively on relevant health outcomes to provide safe and cost-effective care. Efforts to standardise treatment options and reduce selective reporting of outcomes are warranted to reduce inconsistency in evidence synthesis.

Conclusions

430 Medical treatments for first-trimester miscarriage have similar effectiveness and side effects
431 compared to surgery. The addition of MIFE could increase the effectiveness of MISO and
432 reduce side effects though evidence is limited due to inconsistency. EXP has lower
433 effectiveness compared to other treatment options.

434

435 **Acknowledgements:**

436 The authors acknowledge the initial contribution of Arri Coomarasamy, Ioannis Gallos and
437 Mary Eyo.

438

439 **Authors' roles:**

440 BHA conceived the idea, performed the search, extracted data and wrote the first draft. NM
441 extracted data; AT and JZ performed the analysis and revised the manuscript; KSK revised
442 the manuscript and supervised the study. All authors provided critical input to the final
443 manuscript.

444

445 **Funding:**

446 No funding was received in support of this work.

447

448

449 **Conflict of interest:**

450 None

451

452

453 **Figures and tables legends:**

Figure 1: The study selection process for network meta-analysis on management of first trimester miscarriage.

Figure 2: Network of treatment options for first trimester miscarriage. Options: expectant management (EXP), sharp dilation and curettage (D+C), electric vacuum aspiration (EVAC), manual vacuum aspiration (MVA), misoprostol alone (MISO), mifepristone+misoprostol (MIFE+MISO) or misoprostol+electric vacuum aspiration (MISO+EVAC).

The size of the dots represents the number of women randomised to each treatment option and the thickness of the lines represents the number of randomised trials with head to head comparison between each two treatment options.

Figure 3: Direct (D) and mixed (M) evidence meta-analysis for treatment options for first trimester miscarriage. Options: expectant management (EXP), sharp dilation and curettage (D+C), electric vacuum aspiration (EVAC), manual vacuum aspiration (MVA), misoprostol alone (MISO), mifepristone+misoprostol (MIFE+MISO) or misoprostol+electric vacuum aspiration (MISO+EVAC).

Figure (4): The mean rank and cumulative rank probability (SUCRA) of effectiveness for each treatment option for first trimester miscarriage. Options: expectant management (EXP), sharp dilation and curettage (D+C), electric vacuum aspiration (EVAC), manual vacuum aspiration (MVA), misoprostol alone (MISO), mifepristone+misoprostol (MIFE+MISO) or misoprostol+electric vacuum aspiration (MISO+EVAC).

Treatments with the top mean rank and the largest area under the curve have the highest probability of achieving the primary outcome of complete evacuation of products of conception.

Table I: Characteristics of included trials evaluating treatment options for first trimester miscarriage.

Table II: Summary of the calculated mean rank and the surface under the cumulative ranking curve (SUCRA) for the secondary outcomes for the treatment options for first trimester miscarriage.

Treatments ranked first have lower likelihood to achieving adverse outcomes and higher likelihood of post-treatment satisfaction. Treatments with a higher SUCRA score have lower likelihood of achieving adverse outcomes and higher likelihood of post-treatment satisfaction.

Supplementary Figure S1: Risk of bias in included trials on the treatment options for first trimester miscarriage.

Supplementary Figure S2: Flow chart for the management of women with first trimester miscarriage.

Supplementary Figure S3: Mixed evidence meta-analysis adjusted for inconsistency for treatment options for first trimester miscarriage.

Supplementary Figure S4: Funnel plot of the treatment effect for included trials on treatment options for first trimester miscarriage.

Supplementary Figure S5: Mixed evidence network meta-analysis of blood transfusion following treatment options for first trimester miscarriage. (A) Network map. (B) Forest plot. (C) SUCRA.

Supplementary Figure S6: Mixed evidence network meta-analysis of infection/pelvic inflammatory disease following treatment options for first trimester miscarriage. (A) Network map. (B) Forest plot. (C) SUCRA.

Supplementary Figure S7: Mixed evidence network meta-analysis of serious complications following treatment options for first trimester miscarriage. (A) Network map. (B) Forest plot. (C) SUCRA.

Supplementary Figure S8: Mixed evidence network meta-analysis of diarrhoea following treatment options for first trimester miscarriage. (A) Network map. (B) Forest plot. (C) SUCRA.

Supplementary Figure S9: Mixed evidence network meta-analysis of vomiting following treatment options for first trimester miscarriage. (A) Network map. (B) Forest plot. (C) SUCRA.

Supplementary Figure S10: Mixed evidence network meta-analysis of nausea following treatment options for first trimester miscarriage. (A) Network map. (B) Forest plot. (C) SUCRA.

Supplementary Figure 11: Mixed evidence network meta-analysis of fever following treatment options for first trimester miscarriage. (A) Network map. (B) Forest plot. (C) SUCRA.

Supplementary Figure 12: Mixed evidence network meta-analysis of analgesia following treatment options for first trimester miscarriage. (A) Network map. (B) Forest plot. (C) SUCRA.

Supplementary Figure 13: Mixed evidence network meta-analysis of women's satisfaction following treatment options for first trimester miscarriage. (A) Network map. (B) Forest plot. (C) SUCRA.

Supplementary Table SI: Side-split analysis of inconsistency in the network of treatment options for first trimester miscarriage.

Supplementary Table SII: Summary of risk of bias for included studies.

Supplementary Table SIII: League table of effect estimates for secondary outcomes across treatment options for first trimester miscarriage.

References

- Autry A, Jacobson G, Sandhu R, Isbill K. Medical management of non-viable early first trimester pregnancy. *Int J Gynecol Obstet* 1999;**67**:9–13.
- Bagratee JS, Khullar V, Regan L, Moodley J, Kagoro H. A randomized controlled trial comparing medical and expectant management of first trimester miscarriage. *Hum Reprod* 2004;**19**:266–271.
- van den Berg MMJ, Dancet EAF, Erlikh T, van der Veen F, Goddijn M, Hajenius PJ. Patient-centered early pregnancy care: a systematic review of quantitative and qualitative studies on the perspectives of women and their partners. *Hum Reprod Update* 2017:1–13.
- Blohm F, Friden BE, Milsom I, Platz-Christensen JJ, Nielsen S. A randomised double blind trial comparing misoprostol or placebo in the management of early miscarriage. *BJOG An Int J Obstet Gynaecol* 2005;**112**:1090–1095.
- Bucher HC, Guyatt GH, Griffith LE, Walter SD. The results of direct and indirect treatment comparisons in meta-analysis of randomized controlled trials. *J Clin Epidemiol* 1997;**50**:683–691.
- Chaimani A, Higgins JPT, Mavridis D, Spyridonos P, Salanti G. Graphical tools for network meta-analysis in STATA. *PLoS One* 2013;**8**:e76654.
- Dalton VK, Liang A, Hutton DW, Zochowski MK, Fendrick AM. Beyond usual care: the economic consequences of expanding treatment options in early pregnancy loss. *Am J Obstet Gynecol* 2015;**212**:177-e1.
- Fang A, Chen Q, Zheng W, Li Y, Chen R. Termination of Missed Abortion in A Combined Procedure: A Randomized Controlled Trial. *J Reprod Contracept* 2009;**20**:45–49.

574 Frost M, Condon JT. The psychological sequelae of miscarriage: a critical review of the
575 literature. *Aust N Z J Psychiatry* 1996;**30**:54–62.

576 Herabutya Y, O-Prasertsawat P. Misoprostol in the management of missed abortion. *Int J*
577 *Gynecol Obstet* 1997;**56**:263–266.

578 Higgins JPT, Altman DG, Gøtzsche PC, Jüni P, Moher D, Oxman AD, Savović J, Schulz KF,
579 Weeks L, Sterne JAC. The Cochrane Collaboration's tool for assessing risk of bias in
580 randomised trials. *Bmj* 2011;**343**:d5928.

581 Higgins JPT, Jackson D, Barrett JK, Lu G, Ades AE, White IR. Consistency and
582 inconsistency in network meta-analysis: concepts and models for multi-arm studies. *Res*
583 *Synth Methods* 2012;**3**:98–110.

584 Holanda AAR, Dos Santos H, Barbosa MF, Barreto CFB, Felinto AS, De Araújo IS.
585 Tratamento do abortamento do primeiro trimestre da gestação: curetagem versus
586 aspiração manual a vácuo. *RBGO* 2003;**25**:271–276.

587 Hutton B, Salanti G, Caldwell DM, Chaimani A, Schmid CH, Cameron C, Ioannidis JPA,
588 Straus S, Thorlund K, Jansen JP. The PRISMA extension statement for reporting of
589 systematic reviews incorporating network meta-analyses of health care interventions:
590 checklist and explanations. *Ann Intern Med* 2015;**162**:777–784.

591 Al Inizi SAT, Ezimokhai M. Vaginal misoprostol versus dinoprostone for the management of
592 missed abortion. *Int J Gynecol Obstet* 2003;**83**:73–74.

593 Johnson N, Priestnall M, Marsay T, Ballard P, Watters J. A randomised trial evaluating pain
594 and bleeding after a first trimester miscarriage treated surgically or medically. *Eur J*
595 *Obstet Gynecol Reprod Biol* 1997;**72**:213–215.

596 Jurkovic D, Overton C, Bender-Atik R. Diagnosis and management of first trimester
597 miscarriage. *BMJ* 2013;**346**:f3676.

598 Karlsen, Jørn-Hugo; Hjalmar AS. Utskrapning eller ikke etter spontanabort? *Tidsskr Nor*

599 *Lægeforen* 2001;**24**. Available at: [https://tidsskriftet.no/2001/10/klinikk-og-](https://tidsskriftet.no/2001/10/klinikk-og-forskning/utskrapning-eller-ikke-etter-spontanabort)
600 forskning/utskrapning-eller-ikke-etter-spontanabort.

601 Khan K. The CROWN Initiative: journal editors invite researchers to develop core outcomes
602 in women's health. *BJOG* 2016;**123 Suppl**:103–104.

603 Lawrie A, Penney G, Templeton N. A randomised comparison of oral and vaginal
604 misoprostol for cervical priming before suction termination of pregnancy. *BJOG An Int*
605 *J Obstet Gynaecol* 1996;**103**:1117–1119.

606 Lelaidier C, Baton-Saint-Mleux C, Fernandez H, Bourget P, Frydman R. Mifepristone (RU
607 486) induces embryo expulsion in first trimester non-developing pregnancies: a
608 prospective randomized trial. *Hum Reprod* 1993;**8**:492–495.

609 Lemmers M, Verschoor MAC, Hooker AB, Opmeer BC, Limpens J, Huirne JAF, Ankum
610 WM, Mol BWM. Dilatation and curettage increases the risk of subsequent preterm birth:
611 a systematic review and meta-analysis. *Hum Reprod* 2015;**31**:34–45.

612 Lister MS, Shaffer LET, Bell JG, Lutter KQ, Moorma KH. Randomized, double-blind,
613 placebo-controlled trial of vaginal misoprostol for management of early pregnancy
614 failures. *Am J Obstet Gynecol* 2005;**193**:1338–1343.

615 Magotti RF, Munjinja PG, Lema RS, Ngwalle EK. Cost-effectiveness of managing abortions:
616 manual vacuum aspiration (MVA) compared to evacuation by curettage in Tanzania.
617 *East Afr Med J* 1995;**72**:248–251.

618 Molnar AM, Oliver LM, Geyman JP. Patient preferences for management of first-trimester
619 incomplete spontaneous abortion. *J Am Board Fam Pract* 2000;**13**:333–337.

620 Nanda K, Pelligia A, Grimes D, Lopez L, Nanda G. Expectant care versus surgical treatment
621 for miscarriage. *Cochrane Database Syst Rev* 2006;**2**:CD003518.

622 Neilson JP, Gyte GML, Hickey M, Vazquez JC, Dou L. Medical treatments for incomplete
623 miscarriage. *Cochrane Libr* 2013.

624 Neilson JP, Hickey M, Vazquez JC. Medical treatment for early fetal death (less than 24
625 weeks). *Cochrane Libr* 2006.

626 Ngai SW, Chan YM, Tang OS, Ho PC. Vaginal misoprostol as medical treatment for first
627 trimester spontaneous miscarriage. *Hum Reprod* 2001;**16**:1493–1496.

628 Nielsen S, Hahlin M, Platz-Christensen J. Randomised trial comparing expectant with
629 medical management for first trimester miscarriages. *BJOG An Int J Obstet Gynaecol*
630 1999;**106**:804–807.

631 Pereira PP, LONGO ALM, OLIVEIRA FRC, Armelin AR, Maganha CA, Zugaib M.
632 TRATAMENTO DO ABORTAMENTO INCOMPLET INCOMPLETO POR
633 ASPIRAÇÃO MANUAL OU CURETAGEM. *Rev Assoc Med Bras* 2006;**52**:304–307.

634 Queensland Clinical Guidelines. Early pregnancy loss. 2015. Available at:
635 https://www.health.qld.gov.au/__data/assets/pdf_file/0033/139947/g-epl.pdf.

636 Riley RD, Jackson D, Salanti G, Burke DL, Price M, Kirkham J, White IR. Multivariate and
637 network meta-analysis of multiple outcomes and multiple treatments: rationale,
638 concepts, and examples. *bmj* 2017;**358**:j3932.

639 Royal College of Obstetricians and Gynaecologists. Blood Transfusion in Obstetrics. Green-
640 top Guideline No. 47. 2015. Available at:
641 <https://www.rcog.org.uk/globalassets/documents/guidelines/gtg-47.pdf>.

642 Salanti G, Ades AE, Ioannidis JPA. Graphical methods and numerical summaries for
643 presenting results from multiple-treatment meta-analysis: an overview and tutorial. *J*
644 *Clin Epidemiol* 2011;**64**:163–171.

645 Sinha P, Suneja A, Guleria K, Aggarwal R, Vaid NB. Comparison of Mifepristone Followed
646 by Misoprostol with Misoprostol Alone for Treatment of Early Pregnancy Failure: A
647 Randomized Double-Blind Placebo-Controlled Trial. *J Obstet Gynecol India*
648 2018;**68**:39–44.

649 Sotiriadis A, Makrydimas G, Papatheodorou S, Ioannidis JPA. Expectant, medical, or
650 surgical management of first-trimester miscarriage: a meta-analysis. *Obstet Gynecol*
651 2005;**105**:1104–1113.

652 Spitz IM, Bardin CW, Benton L, Robbins A. Early pregnancy termination with mifepristone
653 and misoprostol in the United States. *N Engl J Med* 1998;**338**:1241–1247.

654 Strand EA. Increasing the management options for early pregnancy loss: the economics of
655 miscarriage. *Am J Obstet Gynecol* 2015;**212**:125–126.

656 Sutton AJ, Abrams KR, Jones DR, Jones DR, Sheldon TA, Song F. Methods for meta-
657 analysis in medical research. 2000.

658 Swanson KM, Chen H-T, Graham JC, Wojnar DM, Petras A. Resolution of depression and
659 grief during the first year after miscarriage: a randomized controlled clinical trial of
660 couples-focused interventions. *J Women's Heal* 2009;**18**:1245–1257.

661 Tang OS, Gemzell-Danielsson K, Ho PC. Misoprostol: pharmacokinetic profiles, effects on
662 the uterus and side-effects. *Int J Gynaecol Obstet* 2007;**99 Suppl 2**:S160-7.

663 The American College of Obstetricians and Gynecologists. Early pregnancy loss, Practice
664 Bulletin No. 150. 2015;**125**:1258–6. Available at: [http://www.acog.org/-](http://www.acog.org/-/media/Practice-Bulletins/Committee-on-Practice-Bulletins----Gynecology/Public/pb150.pdf?dmc=1&ts=20170408T1003028826)
665 [/media/Practice-Bulletins/Committee-on-Practice-Bulletins----](http://www.acog.org/-/media/Practice-Bulletins/Committee-on-Practice-Bulletins----Gynecology/Public/pb150.pdf?dmc=1&ts=20170408T1003028826)
666 [Gynecology/Public/pb150.pdf?dmc=1&ts=20170408T1003028826](http://www.acog.org/-/media/Practice-Bulletins/Committee-on-Practice-Bulletins----Gynecology/Public/pb150.pdf?dmc=1&ts=20170408T1003028826).

667 The National Institute for Health and Care Excellence. Ectopic pregnancy and miscarriage:
668 diagnosis and initial management. 2012. Available at:
669 <https://www.nice.org.uk/guidance/cg154>.

670 Trinder J, Brocklehurst P, Porter R, Read M, Vyas S, Smith L. Management of miscarriage:
671 expectant, medical, or surgical? Results of randomised controlled trial (miscarriage
672 treatment (MIST) trial). *bmj* 2006;**332**:1235–1240.

673 Tunçalp Ö, Gülmezoglu AM, Souza JP. Surgical procedures for evacuating incomplete

674 miscarriage. *Cochrane Libr* 2010.

675 Veroniki AA, Vasiliadis HS, Higgins JPT, Salanti G. Evaluation of inconsistency in networks
676 of interventions. *Int J Epidemiol* 2013;**42**:332–345.

677 Wallace RR, Goodman S, Freedman LR, Dalton VK, Harris LH. Counseling women with
678 early pregnancy failure: utilizing evidence, preserving preference. *Patient Educ Couns*
679 2010;**81**:454–461.

680 Wang X, Chen C, Wang L, Chen D, Guang W, French J. Conception, early pregnancy loss,
681 and time to clinical pregnancy: a population-based prospective study. *Fertil Steril*
682 2003;**79**:577–584.

683 Al Wattar BH, Zamora J, Khan KS. Informing treatment decisions through meta-analysis: to
684 network or not? *Evid Based Med* 2017;**22**:12–15.

685 Wen J, Cai QY, Deng F, Li YP. Manual versus electric vacuum aspiration for first-trimester
686 abortion: a systematic review. *BJOG An Int J Obstet Gynaecol* 2008;**115**:5–13.

687 White IR. Multivariate random-effects meta-regression: updates to mvmeta. *Stata J*
688 2011;**11**:255.

689 White IR, Barrett JK, Jackson D, Higgins J. Consistency and inconsistency in network
690 meta-analysis: model estimation using multivariate meta-regression. *Res Synth Methods*
691 2012;**3**:111–125.

692 Wood SL, Brain PH. Medical management of missed abortion: a randomized clinical trial.
693 *Obstet Gynecol* 2002;**99**:563–566.

694